



Australian
National
University

Lithospheric structure in the vicinity of the 11GA-YO1 (YOM) reflection profile

B.L.N. Kennett


Research School of Earth Sciences

The Australian National University



Seismic coverage for Western Central Australia

- The YOM (Yilgarn-Officer-Musgrave) project is part of a extensive network of seismic information
- Prior refraction work and extensive broad-band deployments (Receiver Functions) provide control additional to the reflection work





AuSREM


- The Australian Seismological Reference Model (AuSREM) is an effort to bring together existing information on the 3-D structure beneath the Australian area to create a **representative** model
- Depends on collaborative support from a wide range of researchers, whose work bears on the structure of Australia
- Published Products :
 - Revised Moho Map of Australia – Kennett, Salmon et al. 2011,
 - Updated 2012 Salmon et al, 2013, Tectonophysics – in press
 - Summary Paper – Kennett & Salmon, 2012 AJES
 - Crustal & Mantle Papers 2013, GJI, vol 192.

- Prior compilations of sediment thicknesses
- P and S wavespeed distributions through the crust.
- The primary information for P wavespeed comes from refraction profiles, for S wavespeed from receiver function studies.
- Ambient noise tomography for S wavespeed links the point observations into national coverage.
- Density values are derived using results from gravity interpretations.
- Updated map of depth to Moho, using latest full crustal reflection profiling




Construction of Crustal component


1. Refraction + receiver function V_p field (onshore & offshore)
2. V_p/V_s field from smoothed receiver functions
3. V_p for continent from ambient noise by conversion of V_s
4. Construct composite P wavespeed field from all P results
5. Create S wavespeed from P using V_p/V_s field
6. Construct density from P wavespeed




V_p/V_s
10 km




V_p/V_s
35 km




Construction of Vp field from ambient noise results






Construction of Vp field






Crustal Structure near YOM






Crustal Structure near YOM





Crustal Structure near YOM






Comparison of Moho Depth estimates

The map summarises all the estimates for Moho depth in the neighbourhood of the YOM line

Triangles denote depth estimates from reflection work and refraction.

Diamonds, pentagons and squares represent results from Receiver Functions.

The YOM results are the main constraint for a large area.






Moho variation across Western to Central Australia

Portion of the 2012 Moho map for Australia (Kennett et al. 2011., GJI, updated)


The map is rendered using 0.5x0.5 deg pixels





Moho depth superimposed on tectonic framework


The correlation of thinner crust with the ancient Archean blocks is very clear, as also is thickening at the craton margins.






Passive Seismology – Mantle Studies

- A major tool for studying the mantle lithosphere is the use of surface wave tomography exploiting the large amplitude surface waves generated by shallow earthquakes
- Waveforms are matched with calculations to extract information on structure and build 3-D models
- Can also use higher frequency body wave arrivals



Broad-band stations and regional events (one year)






AuSREM: Mantle Component

- Primary source of information comes from seismic surface wave tomography (multiple studies)
- Representative model has been developed to capture the features of a range of studies for SV and SH
- Body wave studies and regional tomography provide useful constraints on the relation between P and S wavespeeds.
- The mantle model extends beyond the continent and so covers a larger area than for the crust.
- Below 300 km and in the surrounding area AuSREM is linked to the S40RTS model (Ritsema et al., 2011).




SV wavespeed





Base of Lithosphere



Depth to base of lithosphere based on wavespeed gradients

Acknowledgments

- Surface Wave Tomography: S. Fishwick, K.Yoshizawa, A. Fichtner
- Receiver Functions (WA): A. Reading
- Ambient Noise Tomography: E. Saygin
- AuSREM Crust & Moho map compilation: M. Salmon

- All the members of RSES who have helped with the collection of portable seismic data and subsequent data handling.